

Zeros of Witten zeta functions and absolute limit

N. Kurokawa and H. Ochiai

1 Introduction

The Witten zeta function

$$\zeta_G^W(s) = \sum_{\rho \in \hat{G}} \deg(\rho)^{-s}$$

was introduced by Witten [W] in 1991, where G is a compact topological group and $\hat{G}$ denotes the unitary dual, that is, the set of equivalence classes of irreducible unitary representations. The example

$$\zeta_{SU(2)}^W(s) = \sum_{m=0}^{\infty} \deg(\text{Sym}^m)^{-s} = \sum_{n=1}^{\infty} n^{-s} = \zeta(s),$$

where $\zeta(s)$ denotes the Riemann zeta function, suggests fine properties for general case. In fact, Witten showed arithmetical interpretation for $\zeta_{SU(n)}^W(2m)$ ($m = 1, 2, 3, \dots$) containing Euler's result ([E1] 1735)

$$\zeta_{SU(2)}^W(2m) \in \pi^{2m} \mathbf{Q}.$$

In this paper we look at the opposite side: special values at negative integers such as

$$\zeta_{SU(2)}^W(-1) = \left(\sum_{n=1}^{\infty} n \right) = -\frac{1}{12}, \quad (1)$$

$$\zeta_{SU(2)}^W(-2) = \left(\sum_{n=1}^{\infty} n^2 \right) = 0 \quad (2)$$

due to Euler [E2](1749). We notice that the value

$$\left(\sum_{n=1}^{\infty} n \right) = -\frac{1}{12}$$

appears as the one-dimensional Casimir energy: see Casimir [C] and Hawking [H]. The equality

$$\text{“} \sum_{n=1}^{\infty} n^2 \text{”} = 0$$

means the vanishing of the two-dimensional Casimir energy.

We notice that

$$\zeta_G^W(-2) = |G|$$

when G is a finite group. We conjecture that

$$\zeta_G^W(-2) = 0 \tag{3}$$

for infinite groups G .

For deeper understanding of the situation, we introduce a new zeta function (Witten L -function)

$$\zeta_G^W(s, g) = \sum_{\rho \in \widehat{G}} \frac{\text{trace}(\rho(g))}{\deg(\rho)} \deg(\rho)^{-s} \tag{4}$$

where G is a compact topological group, g is an element of G , $\widehat{G}$ is the set of equivalence classes of irreducible ($\mathbf{C}$ -valued) representations of G , $\deg(\rho)$ is the degree (the dimension) of an irreducible representation $\rho \in \widehat{G}$. Note that $\text{trace}(\rho(g))$ is the character of the representation ρ . This Witten zeta function $\zeta_G^W(s, g)$ reduces to the (usual) Witten zeta function when we specialize g to be the identity element $1 \in G$:

$$\zeta_G^W(s) = \zeta_G^W(s, 1).$$

In the case of a finite group G we have

$$\zeta_G^W(-2, g) = \begin{cases} |G| & \text{if } g = 1, \\ 0 & \text{otherwise.} \end{cases}$$

We conjecture that

$$\zeta_G^W(-2, g) = 0 \tag{5}$$

when G is an infinite group. The following result treats the case $G = SU(2)$.

Theorem 1. *Suppose $g \in SU(2)$ is conjugate to $\begin{pmatrix} e^{i\theta} & 0 \\ 0 & e^{-i\theta} \end{pmatrix}$ with $0 \leq \theta \leq \pi$.*

(1) We have an expression

$$\zeta_{SU(2)}^W(s, g) = \sum_{n=1}^{\infty} \frac{\sin(n\theta)}{n \sin \theta} n^{-s}$$

in $\text{Re}(s) > 1$ The function $\zeta_{SU(2)}^W(s, g)$ in s has a meromorphic continuation to the whole complex plane.

(2) For a positive even integer m , we have $\zeta_{SU(2)}^W(-m, g) = 0$ for all $g \in SU(2)$. Moreover, $s = -2$ is a simple zero of $\zeta_{SU(2)}^W(s, g)$, and the first derivative at $s = -2$ is given as

$$\frac{\partial \zeta_{SU(2)}^W}{\partial s}(-2, g) = \begin{cases} -\frac{\zeta(3)}{4\pi^2} & \text{if } \theta = 0, \\ \frac{1}{4\pi \sin \theta} \left(\zeta\left(2, \frac{\theta}{2\pi}\right) - \frac{\pi^2}{2 \sin^2 \frac{\theta}{2}} \right) > 0 & \text{if } 0 < \theta < \pi, \\ \frac{7\zeta(3)}{4\pi^2} & \text{if } \theta = \pi. \end{cases}$$

Here $\zeta(s, x)$ denotes the Hurwitz zeta function.

(3) The special value at $s = -1$ is given as

$$\zeta_{SU(2)}^W(-1, g) = \begin{cases} -\frac{1}{12} & \text{if } \theta = 0, \\ \frac{1}{4 \sin^2 \frac{\theta}{2}} & \text{if } 0 < \theta < \pi, \\ \frac{1}{4} & \text{if } \theta = \pi. \end{cases}$$

We now introduce a ‘multi’-version of Witten L -function. For $g_1, \dots, g_r \in G$, we define

$$\begin{aligned} \zeta_G^W(s; g_1, \dots, g_r) &:= \sum_{\rho \in \widehat{G}} \frac{\text{trace}(\rho(g_1))}{\deg(\rho)} \dots \frac{\text{trace}(\rho(g_r))}{\deg(\rho)} \times \deg(\rho)^{-s} \\ &= \sum_{\rho \in \widehat{G}} \frac{\text{trace}(\rho(g_1)) \dots \text{trace}(\rho(g_r))}{\deg(\rho)^{s+r}}. \end{aligned}$$

It is natural to ask whether the vanishing $\zeta_G^W(-2; g_1, \dots, g_r) \stackrel{?}{=} 0$ of the special value at $s = -2$ for this generalization holds. We have a partial answer to this question.

Theorem 2. We have $\zeta_{SU(2)}^W(-m; g_1, g_2) = 0$ for $g_1, g_2 \in SU(2)$, and a positive even integer m .

We also give an example of the non-vanishing for the case $r = 3$: for some $g \in SU(2)$, we prove that $\zeta_{SU(2)}^W(-2; g, g, g) \neq 0$. These results related with the Lie group $SU(2)$ are given in Section 2.

We report further examples of zeros of Witten zeta functions for infinite groups.

Theorem 3. $\zeta_{SU(3)}^W(s) = 0$ for $s = -1, -2, \dots$.

The proof of this theorem is given in Section 3.

The next example is not a Lie group, but a totally disconnected group. Let $\mathbf{Z}_p$ be the p -adic integer ring for a prime number p .

Theorem 4. Suppose $p \neq 2$. Then $\zeta_{SL_2(\mathbf{Z}_p)}^W(s) = 0$ for $s = -1, -2$.

Now we consider the congruence subgroups. For a positive integer m , we define a subgroup of $SL_3(\mathbf{Z}_p)$ of finite index by

$$SL_3(\mathbf{Z}_p)[p^m] = \ker(SL_3(\mathbf{Z}_p) \rightarrow SL_3(\mathbf{Z}_p/(p^m))).$$

Theorem 5. Suppose $p \neq 3$.

(1)

$$\begin{aligned} \zeta_{SL_3(\mathbf{Z}_p)[p^m]}^W(s) &= p^{8m} \frac{(1 - p^{-2-s})(1 - p^{-1-s})}{(1 - p^{1-2s})(1 - p^{2-3s})} \\ &\quad \times (1 + (p^{-1} + p^{-2})p^{-s} + (1 + p^{-1})p^{-2s} + p^{-2-3s}). \end{aligned}$$

(2) $\zeta_{SL_3(\mathbf{Z}_p)[p^m]}^W(s) = 0$ for $s = -1, -2$.

(3)

$$\zeta_{SL_3(\mathbf{Z}_1)[1^m]}^W(s) = \frac{(s+1)(s+2)}{(s-\frac{1}{2})(s-\frac{2}{3})}.$$

Here we interpret that if $\zeta_{SL_3(\mathbf{Z}_p)[p^m]}^W(s)$ has an expression as an analytic function on p , and there is a limit $p \rightarrow 1$, then its limit is denoted by

$$\zeta_{SL_3(\mathbf{Z}_1)[1^m]}^W(s) = \lim_{p \rightarrow 1} \zeta_{SL_3(\mathbf{Z}_p)[p^m]}^W(s).$$

These results on totally disconnected groups are given in Section 4.

2 $SU(2)$

2.1 Parametrization of irreducible representations of $SU(2)$

The set of equivalence classes, $\widehat{G}$, of irreducible unitary representations of $G = SU(2)$ is parametrized by the set of natural numbers. For a natural number, we denote by $\rho = \rho_n \in \widehat{G}$, the corresponding irreducible representation of G . For a $g = \begin{pmatrix} e^{i\theta} & 0 \\ 0 & e^{-i\theta} \end{pmatrix} \in G$, we have the character formula

$$\text{trace}(\rho(g)) = e^{i(n-1)\theta} + e^{i(n-3)\theta} + \dots + e^{i(3-n)\theta} + e^{i(1-n)\theta} \quad (6)$$

and the degree

$$\deg(\rho) = \text{trace}(\rho(I_2)) = n, \quad (7)$$

where $I_2 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \in SU(2)$ is the identity matrix. We also see that $\text{trace}(\rho(-I_2)) = (-1)^{n-1}n$.

We start from $g = \pm I_2 \in SU(2)$. In these cases, $\zeta_{SU(2)}^W(s, g)$ is written in terms of Riemann zeta function. We see that $\zeta_{SU(2)}^W(s, I_2) = \zeta(s)$, and

$$\zeta_{SU(2)}^W(s, -I_2) = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^s} = (1 - 2^{1-s})\zeta(s). \quad (8)$$

2.2 Poly-logarithm function

We recall the poly-logarithm

$$Z(s, x) = \sum_{n=1}^{\infty} \frac{x^n}{n^s},$$

which is written also as $\text{Li}_s(x)$ in literature. This series converges if $|x| < 1$ and $s \in \mathbf{C}$, or $|x| = 1$ and $\text{Re}(s) > 1$. In the following, we restrict to the case $|x| = 1$.

Theorem 6. *Suppose $|x| = 1$ and $x \neq 1$. Then $Z(s, x)$ is analytically continued to a holomorphic function on $s \in \mathbf{C}$. Moreover, for every non-negative integer m , the function $Z(-m, x)$ can be expressed by a rational function in x . The first several examples are*

$$Z(0, x) = \frac{x}{1-x}, \quad Z(-1, x) = \frac{x}{(1-x)^2}, \quad Z(-2, x) = \frac{x(1+x)}{(1-x)^3}, \dots$$

Proof. For $\operatorname{Re}(s) > 1$, we have

$$\begin{aligned}
Z(s, x) &= x + \frac{x^2}{2^s} + \sum_{n=3}^{\infty} \frac{x^n}{n^s} \\
&= x + \frac{x^2}{2^s} + \sum_{n=2}^{\infty} \frac{x^{n+1}}{(n+1)^s} \\
&= x + \frac{x^2}{2^s} + \sum_{n=2}^{\infty} x^{n+1} n^{-s} (1 + n^{-1})^{-s} \\
&= x + \frac{x^2}{2^s} + \sum_{n=2}^{\infty} x^{n+1} n^{-s} \sum_{k=0}^{\infty} \binom{-s}{k} n^{-k} \\
&= x + \frac{x^2}{2^s} + x \sum_{k=0}^{\infty} \binom{-s}{k} (Z(s+k, x) - x) \\
&= x + \frac{x^2}{2^s} + x(Z(s, x) - x) + x \sum_{k=1}^{\infty} \binom{-s}{k} (Z(s+k, x) - x).
\end{aligned}$$

This shows

$$(1-x)Z(s, x) = x + x^2(2^{-s} - 1) + x \sum_{k=1}^{\infty} \binom{-s}{k} (Z(s+k, x) - x). \quad (9)$$

By the estimates of binomial coefficients, the right-hand side converges absolutely on the right-half plane $\operatorname{Re}(s) > 0$. This shows the analytic continuation of $Z(s, x)$ to $\operatorname{Re}(s) > 0$. Repeating this argument, we obtain the analytic continuation to whole $s \in \mathbf{C}$. To substitute $s = -m$ with $m = 0, 1, \dots$, we have the recursion equation

$$(1-x)Z(-m, x) = x + x^2(2^m - 1) + x \sum_{k=1}^m \binom{m}{k} (Z(-(m-k), x) - x). \quad (10)$$

□

First several examples show

$$\begin{aligned}
Z(-3, x) &= \frac{x(1+4x+x^2)}{(1-x)^4}, & Z(-4, x) &= \frac{x(1+x)(1+10x+x^2)}{(1-x)^5}, \\
Z(-5, x) &= \frac{x(1+26x+66x^2+26x^3+x^4)}{(1-x)^6}.
\end{aligned}$$

These examples seem to show

Lemma 7. Suppose $|x| = 1$ with $x \neq 1$. Then

$$Z(0, x) + Z(0, x^{-1}) = -1, \quad (11)$$

and for every positive integer m ,

$$Z(-m, x) + (-1)^m Z(-m, x^{-1}) = 0. \quad (12)$$

Proof. We start from [Jonquière 1880]

$$e^{-\pi i s/2} Z(s, e^{i\theta}) + e^{\pi i s/2} Z(s, e^{-i\theta}) = \frac{(2\pi)^s}{\Gamma(s)} \zeta(1-s, \frac{\theta}{2\pi}) \quad (13)$$

in Milnor [M]. Putting $s = -m$ with $m = 1, 2, \dots$, we have

$$e^{\pi i m/2} Z(-m, e^{i\theta}) + e^{-\pi i m/2} Z(-m, e^{-i\theta}) = 0.$$

□

We remark that $Z(0, 1) = \zeta(0) = -1/2$. In this sense, the formula (11) is valid also for $x = 1$.

2.3 An example

$$Z(-1, e^{i\theta}) = \frac{1}{(e^{-i\theta/2} - e^{i\theta/2})^2} = -\frac{1}{4 \sin^2(\theta/2)}. \quad (14)$$

and this shows

$$\text{Li}_{-1}(e^{-i\theta}) = \text{Li}_{-1}(e^{i\theta}), \quad (15)$$

an even function in θ .

2.4 Proof of Theorem 1(1) and analytic continuation

Now we consider regular elements in $SU(2)$. Suppose $0 < \theta < \pi$. Then we have, for $\text{Re}(s) > 1$,

$$\begin{aligned} \zeta_{SU(2)}^W \left(s, \begin{pmatrix} e^{i\theta} & 0 \\ 0 & e^{-i\theta} \end{pmatrix} \right) &= \sum_{n=1}^{\infty} \frac{e^{in\theta} - e^{-in\theta}}{e^{i\theta} - e^{-i\theta}} \frac{1}{n} n^{-s} \\ &= \frac{1}{e^{i\theta} - e^{-i\theta}} \sum_{n=1}^{\infty} \left(\frac{e^{in\theta}}{n^{s+1}} - \frac{e^{-in\theta}}{n^{s+1}} \right) \\ &= \frac{1}{e^{i\theta} - e^{-i\theta}} \left\{ Z(s+1, e^{i\theta}) - Z(s+1, e^{-i\theta}) \right\} \\ &= \frac{1}{2i \sin \theta} \left\{ Z(s+1, e^{i\theta}) - Z(s+1, e^{-i\theta}) \right\}, \end{aligned}$$

and the right-hand side has meromorphic continuation to whole $s \in \mathbf{C}$.

Note that we interpret

$$\frac{\sin(n\theta)}{n \sin \theta} = \begin{cases} 1 & \text{if } \theta = 0, \\ (-1)^{n-1} & \text{if } \theta = \pi. \end{cases} \quad (16)$$

2.5 Proof of Theorem 1(2); vanishing

For $g = \pm I_2$ and for positive even integer m , we obtain $\zeta_{SU(2)}^W(-m, \pm I_2) = 0$ from $\zeta(-m) = 0$.

For $g \neq \pm I_2$, suppose $0 < \theta < \pi$. Then for a positive integer m , we have

$$\zeta_{SU(2)}^W(-m, g) = \frac{1}{2i \sin \theta} \left(Z(1 - m, e^{i\theta}) - Z(1 - m, e^{-i\theta}) \right). \quad (17)$$

This is zero for even m by the formula (12).

2.6 Proof of Theorem 1(2), first derivative

We see that

$$\frac{1}{\Gamma(s)} = \frac{s(s+1)}{\Gamma(s+2)} \quad (18)$$

shows that

$$\frac{1}{\Gamma(s)} = -(s+1) + O((s+1)^2), \quad (s \rightarrow -1). \quad (19)$$

We again start from the formula (13)

$$e^{-\pi i s/2} Z(s, x) + e^{\pi i s/2} Z(s, x^{-1}) = \frac{(2\pi)^s}{\Gamma(s)} \zeta\left(1 - s, \frac{\theta}{2\pi}\right)$$

with $x = e^{i\theta}$. Taking $\frac{\partial}{\partial s} \Big|_{s=-1}$ in this formula, we have

$$\begin{aligned} & i \frac{\partial Z}{\partial s}(-1, x) + (-i) \frac{\partial Z}{\partial s}(-1, x^{-1}) \\ & + (-\pi i/2)(i)Z(-1, x) + (\pi i/2)(-i)Z(-1, x^{-1}) = (2\pi)^{-1}(-1)\zeta\left(2, \frac{\theta}{2\pi}\right). \end{aligned}$$

Then

$$i \times 2i \sin \theta \times \frac{\partial \zeta_{SU(2)}^W}{\partial s}(-2, g) = -\pi Z(-1, e^{i\theta}) - \frac{1}{2\pi} \zeta\left(2, \frac{\theta}{2\pi}\right), \quad (20)$$

and

$$4\pi \sin \theta \times \frac{\partial \zeta_{SU(2)}^W}{\partial s}(-2, g) = 2\pi^2 L(-1, e^{i\theta}) + \zeta(2, \frac{\theta}{2\pi}) \quad (21)$$

We have

$$\zeta(2, t) + \zeta(2, 1 - t) = \frac{\pi^2}{\sin^2(\pi t)} \quad (22)$$

since the left-hand side is equal to

$$\sum_{n=0}^{\infty} \frac{1}{(n+t)^2} + \sum_{n=0}^{\infty} \frac{1}{(n+1-t)^2} = \sum_{n=-\infty}^{\infty} \frac{1}{(n+t)^2} \quad (23)$$

which is equal to the right-hand side. This shows

$$8\pi \sin \theta \times \frac{\partial \zeta_{SU(2)}^W}{\partial s}(-2, g) = \zeta(2, \frac{\theta}{2\pi}) - \zeta(2, 1 - \frac{\theta}{2\pi}) > 0 \quad (24)$$

since $\frac{\theta}{2\pi} < 1 - \frac{\theta}{2\pi}$.

2.7 Proof of Theorem 1(3)

$$\zeta_{SU(2)}^W(-1, I_2) = \zeta(-1) = -\frac{1}{12} \quad (25)$$

and

$$\begin{aligned} \zeta_{SU(2)}^W \left(-1, \begin{pmatrix} e^{i\theta} & 0 \\ 0 & e^{-i\theta} \end{pmatrix} \right) &= \frac{Z(0, x) - Z(0, x^{-1})}{x - x^{-1}} \\ &= \frac{-x}{(1-x)^2} = \frac{1}{4 \sin^2(\theta/2)}, \end{aligned} \quad (26)$$

where $x = e^{i\theta}$ for all $0 < \theta \leq \pi$.

2.8 An average over the group

Let G be a finite group. The normalized Haar measure dg on G is, by definition,

$$\int_G f(g) dg = \frac{1}{|G|} \sum_{g \in G} f(g). \quad (27)$$

Then we see that, for all $s \in \mathbf{C}$,

$$\int_G \zeta_G^W(s, g) dg = 1, \quad (28)$$

since the left-hand side is equal to

$$= \sum_{\rho \in \widehat{G}} \left(\int_G \text{trace}(\rho(g)) dg \right) \deg(\rho)^{-s-1}, \quad (29)$$

where the average is non-zero only for the trivial representation ρ .

Now we consider the case where G is a compact group which is not necessarily a finite group. Again let dg be the normalized Haar measure of G so that $\int_G dg = 1$. We ask the value

$$\int_G \zeta_G^W(s, g) dg. \quad (30)$$

We can give some example;

$$\int_{SU(2)} \zeta_{SU(2)}^W(-2, g) dg = 0, \quad (31)$$

$$\int_{SU(2)} \zeta_{SU(2)}^W(-1, g) dg = 1. \quad (32)$$

The latter formula is proved by the Weyl integral formula;

$$\int_{SU(2)} \zeta_{SU(2)}^W(-1, g) dg = \int_0^\pi \zeta_{SU(2)}^W(-1, \begin{pmatrix} e^{i\theta} & 0 \\ 0 & e^{-i\theta} \end{pmatrix}) \frac{2}{\pi} \sin^2 \theta \, d\theta = 1. \quad (33)$$

2.9 $r = 2$

We now discuss the properties of a generalization of Witten zeta functions with several characters. We give a proof of Theorem 2.

Proof.

$$\text{trace}(\rho(g_1)) = \frac{x^n - x^{-n}}{x - x^{-1}}, \text{trace}(\rho(g_2)) = \frac{y^n - y^{-n}}{y - y^{-1}}$$

with $x = e^{i\theta_1}$, $y = e^{i\theta_2}$. In the cases $g_2 = \pm I_2$, we have

$$\zeta_{SU(2)}^W(s, g_1, I_2) = \zeta_{SU(2)}^W(s, g_1), \quad (34)$$

$$\zeta_{SU(2)}^W(s, g_1, -I_2) = \zeta_{SU(2)}^W(s, -g_1), \quad (35)$$

Then the problem on the special values is reduced to the case treated in Theorem 1(2).

Now we may suppose $x, y \neq \pm 1$. Then

$$\begin{aligned}
& \zeta_{SU(2)}^W(s, g_1, g_2) \\
&= \frac{1}{(x - x^{-1})(y - y^{-1})} \sum_{n=1}^{\infty} \frac{(xy)^n + (x^{-1}y^{-1})^n - (xy^{-1})^n - (x^{-1}y)^n}{n^{s+2}} \quad (36) \\
&= \frac{Z(s+2, xy) + Z(s+2, x^{-1}y^{-1}) - Z(s+2, xy^{-1}) - Z(s+2, x^{-1}y)}{(x - x^{-1})(y - y^{-1})}.
\end{aligned}$$

This shows

$$\begin{aligned}
\zeta_{SU(2)}^W(-2, g_1, g_2) &= \frac{(Z(0, xy) + Z(0, x^{-1}y^{-1})) - (Z(0, xy^{-1}) + Z(0, x^{-1}y))}{(x - x^{-1})(y - y^{-1})} \\
&= 0, \quad (37)
\end{aligned}$$

where we have used the formula (11). $\square$

2.10 $r = 3$

By the similar computation, we obtain

$$\begin{aligned}
& \zeta_{SU(2)}^W(s; g, g, g) \\
&= \frac{Z(s+3, x^3) - 3Z(s+3, x) + 3Z(s+3, x^{-1}) - Z(s+3, x^{-3})}{(x - x^{-1})^3}. \quad (38)
\end{aligned}$$

If $x = i$, then

$$\zeta_{SU(2)}^W(-2; g, g, g) = \frac{4Z(1, -i) - 4Z(1, i)}{(2i)^3} = \frac{-2\pi i}{-8i} = \frac{\pi}{4} \neq 0.$$

3 $SU(3)$

3.1 On analytic continuation

Let G be a compact semisimple Lie group. Then the Witten zeta $\zeta_G^W(s)$ has a meromorphic continuation to $\mathbf{C}$. This is a special case of

$$\sum_{m_1, \dots, m_r \geq 1} Q(m_1, \dots, m_r) P(m_1, \dots, m_r)^{-s}. \quad (39)$$

Analytic continuation of these zeta functions is discussed in [Mellin 1900], [Mahler 1928].

3.2 A special value at a negative integer

Let n be a positive integer. Let $M = 2n+2$, and suppose $\operatorname{Re}(s) > -n - \frac{1}{2} + \frac{\varepsilon}{2}$, with $\varepsilon > 0$. By [Ma], we have

$$\begin{aligned}\zeta_{SU(3)}^W(s) &= 2^s \sum_{m,n \geq 1} \frac{1}{m^s n^s (m+n)^s} \\ &= 2^s \frac{\Gamma(2s-1)\Gamma(1-s)}{\Gamma(s)} \zeta(3s-1) \\ &\quad + 2^s \sum_{k=0}^{M-1} (-1)^k \frac{s(s+1) \cdots (s+k-1)}{k!} \zeta(2s+k) \zeta(s-k) \\ &\quad + 2^s \frac{1}{2\pi\sqrt{-1}} \int_{\operatorname{Re}(z)=2n+2-\varepsilon} \frac{\Gamma(s+z)\Gamma(-z)}{\Gamma(s)} \zeta(2s+z) \zeta(s-z) dz.\end{aligned}\tag{40}$$

Reminding

$$\left. \frac{\Gamma(2s-1)}{\Gamma(s)} \right|_{s=-n} = (-1)^{n-1} \frac{n!}{2(2n+1)!},\tag{41}$$

we can put $s = -n$ in this identity and obtain

$$\begin{aligned}\zeta_{SU(3)}^W(-n) &= 2^{-n} (-1)^{n-1} \frac{n!n!}{2(2n+1)!} \zeta(-3n-1) \\ &\quad + 2^{-n} \sum_{k=0}^{2n} (-1)^k \frac{(-n)(1-n) \cdots (k-1-n)}{k!} \zeta(-2n+k) \zeta(-n-k) \\ &\quad + 2^{-n} (-1) \frac{(-n)(1-n) \cdots (-1) \cdot 1 \cdots n}{(2n+1)!} \frac{1}{2} \zeta(-3n-1).\end{aligned}\tag{42}$$

This shows $\zeta_{SU(3)}^W(-n) = 0$ for a positive odd integer n , since $\zeta(-3n-1) = 0$ and $\zeta(-2n+k)\zeta(-n-k) = 0$ for $k = 0, 1, \dots, n$. On the other hand, for a positive even integer n , we have

$$\begin{aligned}\zeta_{SU(3)}^W(-n) &= -2^{-n} \frac{(n!)^2}{(2n+1)!} \zeta(-3n-1) \\ &\quad + 2^{-n} \sum_{k=0}^n \binom{n}{k} \zeta(-2n+k) \zeta(-n-k) = 0,\end{aligned}\tag{43}$$

where the last equality follows from the following lemma:

Lemma 8. *For a positive even integer n , we have*

$$\sum_{k+l=n, k, l \geq 0} \frac{1}{k!l!} \zeta(-n-k) \zeta(-n-l) = \frac{n!}{(2n+1)!} \zeta(-3n-1).\tag{44}$$

Equivalently,

$$\sum_{k+l=n, k, l \geq 0} \frac{1}{k!l!} \frac{B_{n+1+k}}{n+1+k} \frac{B_{n+1+l}}{n+1+l} = -\frac{n!}{(2n+1)!} \frac{B_{3n+2}}{3n+2}. \quad (45)$$

This follows from [CW, Theorem 2] when we substitute $\alpha = \gamma = n - 1$ and $\delta = \varepsilon = 1$. $\square$

This concludes the proof of Theorem 3.

4 The groups over $\mathbf{Z}_p$

4.1 SL_2

Let p be an odd prime. We denote by $\mathbf{Z}_p$ the ring of integers in the non-archimedean local field $\mathbf{Q}_p$. Jaikin-Zapirain [J] obtains the following explicit formula:

$$\zeta_{SL_2(\mathbf{Z}_p)}^W(s) = Z_0(s) + Z_\infty(s), \quad (46)$$

with

$$\begin{aligned} Z_0(s) &= \zeta_{SL_2(\mathbf{F}_p)}^W(s) \\ &= 1 + 2 \left(\frac{p-1}{2} \right)^{-s} + 2 \left(\frac{p+1}{2} \right)^{-s} + \frac{p-1}{2} (p-1)^{-s} \\ &\quad + p^{-s} + \frac{p-3}{2} (p+1)^{-s}, \end{aligned} \quad (47)$$

$$\begin{aligned} Z_\infty(s) &= \frac{1}{1-p^{-s+1}} \left(4p \left(\frac{p^2-1}{2} \right)^{-s} + \frac{p^2-1}{2} (p^2-p)^{-s} \right. \\ &\quad \left. + \frac{(p-1)^2}{2} (p^2+p)^{-s} \right). \end{aligned} \quad (48)$$

This deduces

$$Z_0(-2) = p(p^2-1) = |SL_2(\mathbf{F}_p)| = p(p+1)(p-1), \quad (49)$$

$$Z_\infty(-2) = -p(p^2-1), \quad (50)$$

$$Z_0(-1) = p(p+1), \quad (51)$$

$$Z_\infty(-1) = -p(p+1), \quad (52)$$

$$Z_0(0) = p+4, \quad (53)$$

$$Z_\infty(0) = -\frac{4}{p-1} - p - 4. \quad (54)$$

This shows

$$\zeta_{SL_2(\mathbf{Z}_p)}^W(-2) = 0, \quad (55)$$

$$\zeta_{SL_2(\mathbf{Z}_p)}^W(-1) = 0, \quad (56)$$

$$\zeta_{SL_2(\mathbf{Z}_p)}^W(0) = -\frac{4}{p-1}, \quad (57)$$

which concludes the proof of Theorem 4.

4.2 Congruence subgroups of SL_2

In this subsection, we assume that p is an odd prime. By [AKOV], we obtain

$$\zeta_{SL_2(\mathbf{Z}_p)[p^m]}^W(s) = p^{3m+2} \frac{1 - p^{-2-s}}{1 - p^{1-s}}. \quad (58)$$

This shows

$$\zeta_{SL_2(\mathbf{Z}_p)[p^m]}^W(-2) = 0, \quad (59)$$

$$\zeta_{SL_2(\mathbf{Z}_p)[p^m]}^W(-1) = -p^{3m+1}/(p+1). \quad (60)$$

By taking an “absolute limit” $p \rightarrow 1$, we obtain

$$\zeta_{SL_2(\mathbf{Z}_1)[1^m]}^W(s) = \frac{s+2}{s-1}. \quad (61)$$

4.3 Congruence subgroups of SL_3 and SU_3

In this subsection, we assume that p is a prime with $p \neq 3$. By [AKOV], we have

$$\zeta_{SL_3(\mathbf{Z}_p)[p^m]}^W(s) = p^{8m} \frac{1 + u(p)p^{-3-2s} + u(p^{-1})p^{-2-3s} + p^{-5-5s}}{(1 - p^{1-2s})(1 - p^{2-3s})}, \quad (62)$$

where $u(X) = X^3 + X^2 - X - 1 - X^{-1}$. We notice that it can be factorized as

$$\begin{aligned} \zeta_{SL_3(\mathbf{Z}_p)[p^m]}^W(s) &= p^{8m} \frac{(1 - p^{-2-s})(1 - p^{-1-s})}{(1 - p^{1-2s})(1 - p^{2-3s})} \\ &\quad \times (1 + (p^{-1} + p^{-2})p^{-s} + (1 + p^{-1})p^{-2s} + p^{-2-3s}). \end{aligned} \quad (63)$$

We see that

$$\zeta_{SL_3(\mathbf{Z}_p)[p^m]}^W(-2) = \zeta_{SL_3(\mathbf{Z}_p)[p^m]}^W(-1) = 0.$$

The formula (63) shows

$$\lim_{p \rightarrow 1} \zeta_{SL_3(\mathbf{Z}_p)[p^m]}^W(s) = \frac{(s+1)(s+2)}{(s-\frac{1}{2})(s-\frac{2}{3})}, \quad (64)$$

which is considered to be “an absolute Witten zeta function $\zeta_{SL_3(\mathbf{Z}_1)[1^m]}^W(s)$ ”.

Also by [AKOV],

$$\zeta_{SU_3(\mathbf{Z}_p)[p^m]}^W(s) = p^{8m} \frac{1 + u(p)p^{-3-2s} + u(p^{-1})p^{-2-3s} + p^{-5-5s}}{(1-p^{1-2s})(1-p^{2-3s})} \quad (65)$$

$$\begin{aligned} &= p^{8m} \frac{(1-p^{-2-s})(1-p^{-s})(1+p^{-1-s})}{(1-p^{1-2s})(1-p^{2-3s})} \\ &\quad \times (1 + (1-p^{-1} + p^{-2})p^{-s} + p^{-2-2s}), \end{aligned} \quad (66)$$

where $u(X) = -X^3 + X^2 - X + 1 - X^{-1}$. This shows

$$\zeta_{SU_3(\mathbf{Z}_p)[p^m]}^W(-2) = \zeta_{SU_3(\mathbf{Z}_p)[p^m]}^W(0) = 0,$$

while

$$\zeta_{SU_3(\mathbf{Z}_p)[p^m]}^W(-1) = 2p^{8m-2} \frac{p-1}{p^5-1} = 2p^{8m-2} \frac{1}{[5]_p} \quad (67)$$

is non-zero where $[n]_p = \frac{p^n-1}{p-1}$ is a p -analogue of an integer n . This shows

$$\lim_{p \rightarrow 1} \zeta_{SU_3(\mathbf{Z}_p)[p^m]}^W(-1) = \frac{2}{5}. \quad (68)$$

By the formula (66), we have

$$\lim_{p \rightarrow 1} \zeta_{SU_3(\mathbf{Z}_p)[p^m]}^W(s) = \frac{s(s+2)}{(s-\frac{1}{2})(s-\frac{2}{3})}.$$

References

- [AKOV] Nir Avni, Benjamin Klopsch, Uri Onn, and Christopher Voll, On representation zeta functions of groups and a conjecture of Larsen-Lubotzky, C. R. Acad. Sci. Paris, Ser. I, **348** (2010) 363–367.
- [AKOV2] Nir Avni, Benjamin Klopsch, Uri Onn, and Christopher Voll, Representation zeta functions of some compact p -adic analytic groups, arXiv:1011.6533.
- [C] H. B. G. Casimir, On the attraction between two perfectly conducting plates. Proc. Kon. Ned. Acad. Wetenschap **51**, 793–795 (1948).

- [CW] Venchang Chu and Chenying Wang, Convolution formulae for Bernoulli numbers, *Integral Transforms and Special Functions*, **21** no. 6, (2010), 437–457.
- [E1] L. Euler, De summis serierum reciprocarum (written in 1735), *Commentarii academiae scientiarum Petropolitanae* **7** (1740), 123–134; *Opera Omnia: Series 1, Volume 14*, pp. 73–86.
- [E2] L. Euler, Remarques sur un beau rapport entre les séries des puissances tant directes que réciproques (written in 1749), *Memoires de l’academie des sciences de Berlin* **17** (1768) 83–106; *Opera Omnia: Series 1, Volume 15*, pp. 70–90.
- [H] S.W. Hawking, Zeta function regularization of path integrals in curved space time, *Comm. Math. Phys.* **55** (1977), 133–148.
- [J] A. Jaikin-Zapirain, Zeta function of representations of compact p-adic analytic groups. *J. Amer. Math. Soc.* **19** (2006), 91–118.
- [Jo] A. Jonquière, Note sur la série $\sum_{n=1}^{\infty} \frac{x^n}{n^s}$, *Bull. Soc. Math. France* **17** (1889), 142–152.
- [L] M. Larsen, Determining a semisimple group from its representation degrees. *Internat. Math. Res. Notes* 2004 (2004), 1989–2016.
- [LL] M. Larsen and A. Lubotzky, Representation growth of linear groups. *J. Eur. Math. Soc. (JEMS)* **10** (2008), 351–390.
- [Mah] K. Mahler, Über einen Satz von Mellin, *Math. Ann.* **100** (1928), 384–398.
- [Ma] K. Matsumoto, On analytic continuation of various multiple zeta-functions, In “Number Theory for the Millennium II, Proc. of the Millennium Conference on Number Theory”, M. A. Bennett et al. (eds.), A K Peters, 2002, 417–440.
- [Me] H. Mellin, Eine Formel für den Logarithmus transscendenter Funktionen von endlichem Geschlecht, *Acta Soc. Sci. Fenn.*, **29** (1900), no. 4.
- [M] J. Milnor, On polylogarithms, Hurwitz zeta functions, and the Kubert identities, *Enseign. Math. (2)* **29** (1983), no. 3–4, 281–322.
- [W] E. Witten, On quantum gauge theories in two dimensions, *Comm. Math. Phys.* **141** (1991), 153–209.

[Z] D. Zagier, Values of zeta functions and their applications. In First European Congress of Mathematics, Vol. II (Paris, 1992), Progr. Math. **120**, Birkhäuser, Basel 1994, 497–512.

Nobushige KUROKAWA
Department of Mathematics, Tokyo Institute of Technology,
Oh-Okayama, Meguro, Tokyo, 152-8551, Japan.
`kurokawa@math.titech.ac.jp`

Hiroyuki OCHIAI
Faculty of Mathematics, Kyushu University,
Motooka, Fukuoka, 819-0395, Japan.
`ochiai@imi.kyushu-u.ac.jp`